



## Making and using predictions of species distributions to improve survey design

**Payne, Mark**

*Publication date:*  
2015

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Payne, M. (2015). *Making and using predictions of species distributions to improve survey design*. Abstract from ICES Annual Science Conference 2015, Copenhagen, Denmark.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Making and using predictions of species distributions to improve survey design

Mark R. Payne<sup>1†</sup>

Ecosystem monitoring programmes and environmental niche modelling (ENM, also known as species distribution models) have traditionally had an asymmetrical relationship: data generated by surveys is used to parameterise such models and forms the basis for high-impact publications, but there is rarely feedback in the other direction. However, the knowledge of the factors driving the distribution of a species that ENMs encompass can potentially be of great value to improve the design of surveys. When combined with modern advances in oceanographic observations and modelling, and especially the emerging predictability of some marine systems, the potential exists for great advances. Here I describe the development of a distribution forecast system for Mackerel (*Scomber scombrus*) in the North-east Atlantic. The system has significant forecast and hindcast skill of the spatial distribution of this species, and can therefore be used in real-time (nowcasts) to adapt the design of a survey dynamically. The model also allows prediction of the onset of spawning, thereby allowing the start of the survey to be set to ensure full temporal coverage. Furthermore, the model also shows appreciable forecast skill when coupled to oceanographic forecast models. The application of this approach to the design and execution of the upcoming 2016 Mackerel Egg Survey (MEGS) will be discussed.

Keywords: Environmental niche modelling; Species distribution modelling; Mackerel; predictions; MEGS

<sup>1</sup> National Institute of Aquatic Resources (DTU-Aqua), Technical University of Denmark, 2920 Charlottenlund, Denmark.

<sup>†</sup> Corresponding author. Address: National Institute of Aquatic Resources (DTU-Aqua), Technical University of Denmark, 2920 Charlottenlund, Denmark. Tel.: +45 3396 3455. E-mail address: [mpa@aqu.dtu.dk](mailto:mpa@aqu.dtu.dk)